

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COO

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number
WO 01/92253 A2

- (51) International Patent Classification⁷: **C07D 401/04, A61K 31/4166, A61P 29/00**
- (74) Agent: **MURPHY, Gerald, M., Jr.**; Birch, Stewart, Kolasch & Birch, LLP, P.O. Box 747, Falls Church, VA 22040-0747 (US).
- (21) International Application Number: **PCT/US01/12323**
- (22) International Filing Date: **17 April 2001 (17.04.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/207,968 **31 May 2000 (31.05.2000)** **US**
- (71) Applicant (*for all designated States except US*): **TAN-ABE SEIYAKU CO., LTD.** [JP/JP]; 2-10 Doshomachi 3-chome, Chuo-ku, Osaka 541-8505 (JP).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **SIRCAR, Ila** [US/US]; 4832 Riding Ridge Road, San Diego, CA 92130 (US). **XIE, Yun Feng** [US/US]; 4526 Vereda mar De Ponderosa, San Diego, CA 92130 (US). **SMITH, Nicholas** [US/US]; 1053 Sapphire Street, #1, San Diego, CA 92109 (US). **FURTH, Paul, S.** [US/US]; 5223 Aleman Place, Bonita, CA 91902 (US).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **INHIBITORS OF ALPHA L BETA 2 MEDIATED CELL ADHESION**

(57) Abstract: The present invention relates to small molecules according to formula (I) that are potent inhibitors of $\alpha_L\beta_2$ mediated cell adhesion and which could be useful for the treatment of inflammatory diseases:

WO 01/92253 A2

INHIBITORS OF $\alpha_L\beta_2$ MEDIATED CELL ADHESIONBACKGROUND OF THE INVENTION5 Field of the Invention

The present invention relates to small molecules that are potent inhibitors of $\alpha_L\beta_2$ mediated cell adhesion which could be useful for the treatment of inflammatory diseases.

10 Description of Related Art

The integrin family of proteins are heterodimeric receptors which are expressed on all cell types to mediate cell to cell binding and adhesion to extracellular matrix. The β_2 (CD18) integrin subfamily is comprised of 3 members, 15 $\alpha_L\beta_2$ integrin (LFA-1, CD11a/CD18), $\alpha_M\beta_2$ integrin (Mac-1, CD11b/CD18), and gp 150 β_2 integrin ($\alpha_X\beta_2$ integrin, CD11c/CD18) that are primarily expressed on leukocytes (Sanchez-Madrid et al., J. Exp. Med., 158, 1785-1803 (1983)). $\alpha_L\beta_2$ integrin is found mostly on T and B lymphocytes, while $\alpha_M\beta_2$ integrin is 20 present on activated neutrophils, NK cells and some myeloid cells. The $\alpha_L\beta_2$ integrin binds to intracellular adhesion molecules ICAM-1, 2 and 3 found on multiple cell types such as vascular endothelial cells, dendritic cells, epithelial cells, macrophage and T lymphoblasts (Dustin et al., J. Immunology, 25 137, 245-254 (1986)). Recently there has been evidence presented that $\alpha_L\beta_2$ integrin binds to ICAM-4 and a novel ligand expressed in brain telencephalin. It has been shown that the I domain of the alpha chain is the major recognition site for its ligands.

30 $\alpha_L\beta_2$ integrin adhesion to ICAM-1 is necessary for immune responsiveness of T-lymphocytes to antigens, lymphocyte homing and circulation, and cell emigration to sites of inflammation

(Springer, Ann. Rev. Physiol., 57, 827 (1995)). A dominant role of $\alpha_L\beta_2$ integrin in mediating inflammatory events is shown in several different animal models of inflammatory diseases in which antibodies to $\alpha_L\beta_2$ integrin or ICAM-1 significantly
5 inhibit development of therapeutic end points (Rothlein et al., Kidney International, 41, 617 (1992); Iigo et al., J. Immunology, 147, 4167 (1991); Bennet et al., J. Pharmacol. and Exp. Therapeutics, 280, 988 (1997)).


Also, β_2 integrin subfamily are thought to play a critical
10 role in several types of inflammatory disease processes by interacting with ICAMs. Support for the importance of β_2 integrin in mediating inflammatory responses has been demonstrated by the evidence that transendothelial migration in vitro is markedly inhibited by monoclonal antibodies
15 against β_2 integrin or ICAM-1 (Smith, Can. J. Physiol. Pharmacol., 71, 76 (1993)). Furthermore, blockade of $\alpha_L\beta_2$ integrin has been shown to inhibit neutrophil influx in almost every system, including skin, peritoneum, synovium, lung, kidney, and heart. As one of the primary ligands for the β_2
20 integrin, it would also be expected that blockade of ICAM-1 would inhibit the inflammatory response (Albelda et al., The FASEB Journal, 8, 504 (1994)).

Moreover, it has been shown that antibodies against $\alpha_L\beta_2$ integrin suppress rejection after transplantation. WO
25 94/04188 discloses the use of monoclonal antibodies directed against $\alpha_L\beta_2$ integrin for all transplantations, including graft vs. host or host vs. graft diseases.

SUMMARY OF THE INVENTION

30

The present invention relates to a compound of the formula (I):



or a pharmaceutically acceptable salt thereof, wherein R^1 is selected from

- 1) hydrogen atom, or
 - 2) a C_{1-6} alkyl group which may be optionally substituted with a carboxyl group or a C_{1-6} alkoxycarbonyl group;
- R^{2a} and R^{2b} are independently hydrogen atom, a halogen atom, hydroxyl group, cyano group, a C_{1-6} alkyl group which may be optionally substituted with 1 to 3 halogen atoms, a C_{1-6} alkylthio group which may be optionally substituted with 1 to 3 halogen atoms, a C_{1-6} alkylsulfinyl group which may be optionally substituted with 1 to 3 halogen atoms, a C_{1-6} alkylsulfonyl group which may be optionally substituted with 1 to 3 halogen atoms, or a C_{1-6} alkoxy group which may be optionally substituted with 1 to 3 halogen atoms;
- R^3 is a C_{1-6} alkyl group; and
- R^4 and R^5 are independently a halogen atom.

The compound of the present invention has potent inhibitory activity against $\alpha_L\beta_2$ mediated cell adhesion, and shows excellent in vivo improvements against the unfavorable conditions caused by $\alpha_L\beta_2$ mediated cell adhesion.

Detailed Description of the Invention

The desired compound of the present invention may exist in the form of optical isomers based on asymmetric atoms

thereof, and the present invention also includes these optical isomers and mixtures thereof.

In an embodiment of the present invention, the steric configuration of a bond need not be fixed. The compound of
5 the present invention may be a compound with a sole configuration or a mixture with several different configurations.

In a preferred embodiment of the compound (I), R^1 is hydrogen atom or a C_{1-6} alkyl group which may be optionally
10 substituted with carboxyl group or a C_{1-6} alkoxy carbonyl group, R^{2a} and R^{2b} are independently hydrogen atom, a halogen atom, hydroxyl group, cyano group or a C_{1-6} alkoxy group which may be optionally substituted with 1-3 halogen atoms, R^3 is a C_{1-6} alkyl group, and R^4 and R^5 are independently a halogen atom.

15 In a more preferred embodiment of the compound (I), R^1 is hydrogen atom or a C_{1-6} alkyl group, one of R^{2a} and R^{2b} is hydrogen atom, and the other is a halogen atom, cyano group, or a C_{1-6} alkoxy group which may be optionally substituted with 1-3 halogen atoms, R^3 is a C_{1-6} alkyl group, and R^4 and R^5 are
20 independently a halogen atom.

In a further preferred embodiment of the compound (I), R^1 is hydrogen atom or methyl group, one of R^{2a} and R^{2b} is hydrogen atom and the other is bromine atom, cyano group, a C_{1-6} alkoxy group or trifluoromethoxy group, R^3 is methyl group, R^4 and R^5
25 are chlorine atom.

In another more preferred embodiment of the compound (I), R^1 is hydrogen atom or C_{1-6} alkyl group which may be substituted with carboxyl or C_{1-6} alkoxy carbonyl, one of R^{2a} and R^{2b} is hydrogen atom, and the other is cyano group or C_{1-6} alkoxy
30 group which may be substituted with 1-3 halogen atoms.

In another further preferred embodiment of the compound (I), R^1 is hydrogen atom or methyl group, one of R^{2a} and R^{2b} is hydrogen atom and the other is C_{1-6} alkoxy group or

trifluoromethoxy group, R³ is methyl group, R⁴ and R⁵ are chlorine atom.

In another preferred embodiment of the compound (I), R¹ is a C₁₋₆ alkyl group which is substituted with a C₁₋₆ alkoxy carbonyl group or carboxyl group, one of R^{2a} and R^{2b} is hydrogen atom, and the other is a halogen atom, cyano group, or C₁₋₆ alkoxy group which may be optionally substituted with 1-3 halogen atoms, R³ is a C₁₋₆ alkyl group, and R⁴ and R⁵ are independently a halogen atom.

10 In a more preferred embodiment of the compound (I), R³ is methyl group, and R⁴ and R⁵ are chlorine atom.

Most preferred compound of the present invention is selected from:

3-(2,6-Dichloro-4-pyridyl)-5-(4-bromobenzyl)-1,5-dimethyl-2,4-
15 imidazolidinedione,

3-(2,6-Dichloro-4-pyridyl)-5-(4-propoxybenzyl)-1,5-dimethyl-
2,4-imidazolidinedione,

3-(2,6-Dichloro-4-pyridyl)-5-(4-ethoxybenzyl)-1,5-dimethyl-
2,4-imidazolidinedione,

20 3-(2,6-Dichloro-4-pyridyl)-5-(4-(1,1,1trifluoromethoxybenzyl))-
-5-methyl-2,4-imidazolidinedione,

3-(2,6-dichloro-4-pyridyl)-5-[4-(1,1,1trifluoromethoxybenzyl)]
-1,5-Dimethyl-2,4-imidazolidinedione,

3-(2,6-Dichloro-4-pyridyl)-5-(4-Cyanobenzyl)-5-methyl-2,4-
25 imidazolidinedione,

3-(2,6-Dichloro-4-pyridyl)-5-(4-Cyanobenzyl)-1,5-dimethyl-2,4-
imidazolidinedione; and

a pharmaceutically acceptable salt of these compounds.

The compound of the present invention has potent
30 inhibitory activity against $\alpha_L\beta_2$ mediated cell adhesion, and also shows excellent bioavailability after oral administration which reflects the overall improvement in plasma protein binding and solubility. The compound of the present invention

therefore shows excellent in vivo improvements against the unfavorable conditions caused by $\alpha_L\beta_2$ mediated cell adhesion.

The compound of the present invention may be clinically used either in a free form or in the form of pharmaceutically acceptable salts thereof. Pharmaceutically acceptable salts include an acid-addition salt with an inorganic acid or an organic acid (e.g., hydrochloride, sulfate, nitrate, hydrobromide, methanesulfonate, p-toluenesulfonate, acetate), and a salt with an inorganic base, an organic base or an amino acid (e.g., triethylamine salt, a salt with lysine, an alkali metal salt, an alkali earth metal salt and the like). Pharmaceutically acceptable salts also include an intramolecular salt thereof, or a solvate or hydrate thereof.

The compound of the present invention may be formulated into a pharmaceutical composition comprising a therapeutically effective amount of the compound as defined above and a pharmaceutically acceptable carrier or diluent. The pharmaceutically acceptable carrier or diluent may be, for example, binders (e.g., syrup, gum arabic, gelatin, sorbitol, tragacanth, polyvinylpyrrolidone), excipients (e.g., lactose, sucrose, corn starch, potassium phosphate, sorbitol, glycine), lubricants (e.g., magnesium stearate, talc, polyethylene glycol, silica) disintegrators (e.g., potato starch), wetting agents (e.g., sodium laurylsulfate), and the like.

The desired compound of the present invention or pharmaceutically acceptable salts thereof may be administered either orally or parenterally, and it may be used as a suitable pharmaceutical preparation. These pharmaceutical preparations may be in the form of a solid preparation such as a tablet, a granule, a capsule, and a powder, or in the form of a liquid preparation such as solution, suspension, and emulsion, when administered orally. When administered parenterally, the pharmaceutical preparation may be in the

form of suppository, an injection preparation or an intravenous drip preparation using distilled water for injection, a physiological salt solution, an aqueous glucose solution, and so on, and an inhalation by a conventional
5 process.

The dose of the desired compound of the present invention or a pharmaceutically acceptable salt thereof varies depending on an administration method, age, sex, body weight, and condition of a patient, but, in general, the daily dose is
10 preferably about 0.1 to 100 mg/kg/day, particularly preferably 1 to 100 mg/kg/day.

The compound of the present invention can be used for treating or preventing $\alpha_L\beta_2$ adhesion mediated conditions in a mammal such as a human.

15 The compound of the present invention may be used for treatment or prevention of numerous inflammatory diseases such as rheumatoid arthritis, asthma, allergy conditions, adult respiratory distress syndrome, AIDS, cardiovascular diseases, thrombosis, harmful platelet aggregation, reocclusion
20 following thrombolysis, reperfusion injury, skin inflammatory diseases (e.g., psoriasis, eczema, contact dermatitis, atopic dermatitis), osteoporosis, osteoarthritis, arteriosclerosis (including atherosclerosis), neoplastic diseases including metastasis of neoplastic or cancerous growth, wound, detaching
25 retina, Type I diabetes, multiple sclerosis, systemic lupus erythematosus (SLE), ophthalmic inflammatory conditions, inflammatory bowel diseases (Crohn's disease and ulcerative colitis), regional enteritis, Sjogren's Syndrome, and other autoimmune diseases.


30 The compound of the present invention may also be used for the rejection (i.e., chronic rejection and acute rejection) after transplantation, including allograft rejection (host vs. graft disease) and graft vs. host disease.

The compound of the present invention may be preferably used for treatment or prevention of psoriasis, rheumatoid arthritis, inflammatory bowel diseases (Crohn's disease, ulcerative colitis), systemic lupus erythematosus, atopic
 5 dermatitis, Sjogren's syndrome, and rejection after transplantation (allograft rejection and graft vs. host disease).


According to the present invention, the desired compound
 10 (I) can be prepared by the following methods:

Method A:

Among the desired compound (I), a compound of the formula
 15 (I-a):



wherein the symbols are the same as defined above, or a pharmaceutically acceptable salt thereof, can be prepared by
 (1) cyclizing the compound of the formula (II):



wherein OR⁶ is a hydroxyl group or a protected hydroxyl group, and the other symbols are the same as defined above, and

(2) converting the resulting cyclized compound into a pharmaceutically acceptable salt thereof by a conventional method, if desired.

When OR⁶ is a protected hydroxyl group, the protecting group can be selected from the conventional protecting groups for a carboxyl group (i.e., a C₁₋₆ alkyl group, benzyl group).

The cyclization can be carried out by a conventional condensation method. For example, the cyclization of the compound (II) can be carried out in the presence of an acid or a base in a suitable solvent.

The acid can be selected from organic acids (i.e., p-toluenesulfonic acid, and trifluoroacetic acid) and inorganic acids (i.e., hydrochloric acid, sulfuric acid, and nitric acid).

The base can be selected from conventional bases such as alkali metal alkoxide (e.g., NaOEt, NaOMe).

The solvent can be selected from any one which does not disturb the cyclization reaction, for example, CH₂Cl₂, THF, DMF, alcohols (methanol, ethanol, etc.) or a mixture thereof. The reaction is carried out at a temperature of 0 °C to boiling point of the solvent, preferably at 50 °C to 100 °C.

The cyclization of the compound (II) is also carried out in the presence of a condensing reagent with or without a base in a suitable solvent or without a solvent. The condensing reagent can be selected from SOCl₂ and conventional condensing reagents which can be used for a peptide synthesis, for example, BOP-Cl, BOP reagent, DCC, EDC or CDI.

The base can be selected from an organic base (e.g., DIEA, DMAP, DBU, Et₃N), an alkali metal hydride (e.g., NaH, LiH), an alkali metal carbonate (e.g., Na₂CO₃, K₂CO₃), an alkali metal hydrogen carbonate (e.g., NaHCO₃, KHCO₃), an alkali metal


amide (e.g., NaNH_2), an alkali metal alkoxide (e.g., NaOMe , KOMe), a C_{1-6} alkyl alkali metal salt (e.g., n-BuLi , t-BuLi), an alkali metal hydroxide (e.g., NaOH , KOH), an alkaline earth metal hydroxide (e.g., Ba(OH)_2), and the like.

5 The solvent can be selected from any one which does not disturb the cyclization reaction, for example, CH_2Cl_2 , THF, DMF or a mixture thereof. The reaction is carried out at a temperature of 0°C to room temperature, preferably at room temperature.

10

Method B:

Among the desired compound (I), a compound of the formula (I-b):



15 wherein R^{11} is a C_{1-6} alkyl group which may be optionally substituted with carboxyl group or a C_{1-6} alkoxy carbonyl group and the other symbols are the same as defined above, or a pharmaceutically acceptable salt thereof, may be prepared by:

- (1) alkylating the compound (I-a),
- 20 (2) hydrolyzing the resulting compound, if necessary, and
- (3) converting the resulting compound into a pharmaceutically acceptable salt thereof by a conventional method, if further desired.

(1) Alkylation Reaction

25 The alkylation reaction can be carried out by reacting the compound (I-a) with the compound of the formula (III):



wherein X is a leaving group and R^{11} is the same as defined above.

The leaving group X can be selected from conventional leaving groups, such as a halogen atom (e.g., chlorine, bromine, iodine) and an alkylsulfonyloxy group or an arylsulfonyloxy group (e.g., methylsulfonyloxy group, p-tolylsulfonyloxy group).

The alkylation reaction can be carried out in the presence of a base in a suitable solvent.

10 The base can be selected from conventional bases such as alkali metal hydride (i.e., NaH, KH), alkali metal alkoxide (i.e., NaOMe, NaOEt) and alkali metal amide (i.e., NaNH_2 , LDA, KHMDS).


The solvent can be selected from any one which does not disturb the condensation reaction, for example, DME, THF, DMF, HMPA or a mixture thereof. The reaction is carried out at a temperature of -78°C to room temperature.

(2) Hydrolysis Reaction


The compound (I-b) wherein R^{11} is a C_{1-6} alkyl group substituted with carboxyl group can be prepared by hydrolyzing the compound (I-b) wherein R^{11} is a C_{1-6} alkyl group substituted with a C_{1-6} alkoxycarbonyl group. The hydrolysis can be carried out by a usual procedure, for example, by treating the compound with a base in a suitable solvent. The base can be selected from conventional inorganic bases such as LiOH, NaOH and KOH. The solvent can be selected from any one which does not disturb the hydrolyzing reaction, for example, THF, MeOH, EtOH, H_2O or a mixture thereof. The reaction can be carried out at a temperature of -78°C to 50°C , preferably at a temperature of 0°C to room temperature.

Method C

Among the desired compound (I), the compound of the formula (I-c):



5 or a pharmaceutically acceptable salt thereof, wherein R^{21} is a C_{1-6} alkoxy group and the other symbols are the same as defined above, can be prepared by alkylating a compound of the formula (I-d):




10 wherein the symbols are the same as defined above, and converting into the pharmaceutically acceptable salt, if desired.

The alkylation reaction can be carried out in a similar manner as described in Method B (1) using a suitable halogenated C_{1-6} alkane (e.g., methyl iodide, benzyl bromide) in the presence of a base (e.g., Et_3N , DIEA, $NaHCO_3$, $KHCO_3$, Na_2CO_3 , K_2CO_3 , $KHCO_3$, $CsCO_3$) at a temperature of 0 °C to 50 °C in an organic solvent (e.g., CH_2Cl_2 , THF, DMF, CH_3CN , toluene).

The compound (I) wherein R^{2a} and/or R^{2b} are hydroxyl groups can be prepared by the demethylation of the compound (I) wherein R^{2a} and/or R^{2b} is methoxy group. The demethylation reaction can be carried out by a conventional method, for example, a treatment with BBr_3 or HBr at a temperature of $-78^\circ C$ to $50^\circ C$ in a suitable solvent (e.g., $AcOH$, water).

The starting compound of the formula (II) can be prepared by the following scheme:

10 Scheme 1.



(In the Scheme 1, the symbols are the same as defined above.)

Step 1: The compound (VII) can be prepared by reacting the compound (VIII) with pivalaldehyde. The reaction can be carried out in the presence or absence of an acid or an acidic salt in a suitable solvent or without a solvent. The acid can be selected from conventional inorganic acid such as HCl , H_2SO_4 . The acidic salt can be selected from a salt of a strong inorganic acid and a weak inorganic base such as $MgSO_4$. The solvent can be selected from any one which does not disturb the reaction, for example, toluene, DME, DMF, THF, CH_2Cl_2 or a

mixture thereof. The reaction can be carried out, for example, at a temperature of 0 °C to room temperature.

Step 2: The compound (IV) can be prepared by 1) reacting the compound (VII) with the compound (VI), and 2) hydrolyzing the
5 resulting compound.

The reaction of the compound (VII) and the compound (VI) can be carried out in the presence of a base in a suitable solvent or without a solvent. The base can be selected from conventional bases such as alkali metal alkoxides (e.g., t-
10 BuOK, MeONa, EtONa) and alkali metal amides (e.g., LDA, NaNH₂). The solvent can be selected from any one which does not disturb the coupling reaction, for example, toluene, DME, DMF, THF, CH₂Cl₂ or a mixture thereof. The reaction can be carried out, for example, at a temperature of -78 °C to 50 °C,
15 preferably at a temperature of -10 °C to 0 °C.

Hydrolysis can be carried out in the presence of an acid in a suitable solvent or without a solvent. The acid can be selected from conventional inorganic acid such as HNO₃, HCl, and H₂SO₄. The solvent can be selected from any one which does
20 not disturb the reaction, for example, toluene, DME, DMF, THF, CH₂Cl₂ or a mixture thereof. The reaction can be carried out, for example, at a temperature of 0 °C to room temperature.

Step 3: The compound (II) can be prepared by reacting the compound (IV) with the compound (V).

25 The reaction can be carried out in the presence or absence of a base in a suitable solvent or without a solvent. The base can be selected from conventional inorganic bases such as K₂CO₃, Na₂CO₃ and NaHCO₃, and conventional organic bases such as pyridine, Et₃N, iPr₂EtN, aniline, and N,N-
30 dimethylaniline. The solvent can be selected from any one which does not disturb the coupling reaction, for example, toluene, DME, DMF, THF, CH₂Cl₂ or a mixture thereof. The coupling reaction can be carried out, for example, at a

temperature of -78 °C to 50 °C, preferably at a temperature of 0 °C to room temperature.

In the present description and the claims, the C₁₋₆ alkyl group means a straight chain or branched chain alkyl group having 1 to 6 carbon atoms, for example, methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, etc., preferably one having 1 to 4 carbon atoms. The C₁₋₆ alkoxy means a straight chain or branched chain alkoxy group having 1 to 6 carbon atoms, for example, methoxy group, ethoxy group, propoxy group, isopropoxy group, butoxy group, isobutyloxy, etc., preferably one having 1 to 4 carbon atoms.

Abbreviations

AcOEt:	Ethyl acetate (= EtOAc)
15 BSA:	Bovine serum albumin
DMF:	Dimethyl formamide
DCM:	Dichloromethane
DIEA:	Diisopropylethylamine
DMSO:	Dimethyl sulfoxide
20 Et:	Ethyl
EtOH:	Ethanol
HBSS:	Hank's balanced salt solution
HMPA:	Hexamethylphosphoramide
HSA:	Human serum albumin
25 KHMDS:	Potassium hexamethyldisilazide (=Potassium bis(trimethylsilyl)amide)
LDA:	Lithium diisopropylamide
Me:	Methyl
MeOH:	Methanol
30 n-Bu:	n-Butyl
Ph:	Phenyl
t-Bu:	tert-butyl
THF:	Tetrahydrofuran

Tf: Trifluoromethanesulfonyl


TFA: Trifluoroacetic acid

The compound of the present invention is exemplified by
5 the following examples but not limited thereby.

Examples

Example 1: 3-(2,6-Dichloro-4-pyridyl)-5-(4-cyanobenzyl)-5-methyl-2,4-imidazolidinedione

10



Step-1. L-Alanine ethyl ester hydrogen chloride salt (15 g) was dissolved in H₂O (60 ml). NEt₃ (10.9 g) was added to the
15 stirring solution. The solution was allowed to stir for 30 minutes at room temperature and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and evaporated to give 9.5 g of L-Alanine ethyl ester. The product was used directly for next step. MS: 118 (MH⁺).

20

Step- 2. L-Alanine ethyl ester from step-1 (9 g) was dissolved in 150 ml anhydrous CH_2Cl_2 . The solution was cooled to 0 °C. MgSO_4 (10.17 g) was added to the solution followed by addition of pivalaldehyde (6.95 g). The reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction mixture was filtered and the filtrate was evaporated to give 11.2 g of N-neopentylidene-L-alanine ethyl ester. The product was used for next step without further purification. MS: 186 (MH^+).

10

Step-3. 4-Cyanobenzyl bromide (4.6 g) was added to a solution of the compound obtained from step 3 (4 g) in anhydrous toluene (40 mL). The resulting mixture was cooled to -10°C. t-BuOK (2.9 g) was added portionwise maintaining the temperature at 0 °C. The reaction mixture was stirred at that temperature for 4 hours. The mixture was partitioned between EtOAc/ H_2O . The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. 1 M HCl was added (40 mL) to the residue and the resulting mixture was allowed to stir overnight. EtOAc was added and the reaction mixture was allowed to stir for 30 minutes. The organic phase was separated and the aqueous layer was extracted with additional EtOAc. The combined organic layers were washed with H_2O . The pH of the combined aqueous solution was adjusted to approximately 8 with solid NaHCO_3 , and the mixture was extracted with EtOAc. The combined organic layers were dried over Na_2SO_4 , filtered and concentrated to yield ethyl 2-amino-2-(4-cyanobenzyl)propanoate that was used directly for the next step. MS: 233 (MH^+).

30


Step-4. 2,6-Dichloro-4-pyridyl isocyanate (1 g) was added to a solution of the compound obtained from step-3 (1.35 g) in anhydrous CH_2Cl_2 (10 mL) maintained at 0 °C. The reaction mixture was allowed to warm to room temperature and stirred

overnight. The mixture was partitioned between EtOAc/H₂O. The combined organic layers were dried over Na₂SO₄, filtered and concentrated to give 2-(4-cyanobenzyl)-5-(2,6-dichloro-4-pyridyl)-2-methylhydantoic acid ethyl ester. The product was used directly for the next step. MS: 421 (MH⁺).

Step-5. NaOEt (0.16 g) was added to a solution of the compound from step-4 (1 g) in anhydrous EtOH (10 mL) at 0 °C. The yellow solution was then warmed to room temperature and stirred for 1 hour. EtOH was evaporated and the residue was partitioned between EtOAc/H₂O. The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The product was purified by flash chromatography on silica gel to give the titled compound. (850 mg). MS: 375 (MH⁺).

15


Example 2: 3-(2,6-dichloro-4-pyridyl)-5-(4-Cyanobenzyl)-1,5-Dimethyl-2,4-imidazolidinedione



The compound obtained in Example 1 (400 mg) and t-BuOK (180 mg) were added to a reaction flask and was then flushed with N₂. The mixture was cooled to 0 °C and THF (10 mL) was added. The reaction mixture was stirred at 0 °C for 20 minutes followed by the addition of MeI (454 mg). The reaction mixture was stirred at 0 °C for 3 hours and finally at room temperature for 1 hour. The mixture was extracted with EtOAc/H₂O. The combined organic layers were dried over Na₂SO₄,


filtered and evaporated. The product was purified by prep TLC to give the titled compound (310 mg) MS: 389 (MH⁺).

Example 3: 3-(2,6-Dichloro-4-pyridyl)-5-(4-Cyanobenzyl)-5-methyl-1-(5-ethoxycarbonylpentyl)-2,4-imidazolidinedione



The compound obtained in Example 1 (154 mg) was taken in 2 mL anhydrous DMF. The solution was cooled to 0 °C and NaH (25 mg, 60% in oil) was added. The resulting mixture was stirred for 20 minutes at 0 °C. Ethyl 6-bromohexanoate (140 mg) was added dropwise and the resulting mixture was stirred at room temperature overnight. The reaction mixture was extracted with EtOAc/H₂O. The combined organic layers were dried over Na₂SO₄, filtered and evaporated. The product was purified by preparative TLC to give the titled compound (180 mg). MS: 517 (MH⁺).

Example 4: (R)-3-(2,6-Dichloro-4-pyridyl)-5-(4-Cyanobenzyl)-5-methyl-2,4-imidazolidinedione



Step-1: α -Methyl-4-cyanophenylalanine ethyl ester 1 was prepared according to the methodology described in WO 98/39303.

5

Step-2. To a solution of the compound obtained above (770 mg) in DCM (5 mL) was added HCl (1M in Et₂O, 7 mL) and the reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was then concentrated *in vacuo* to give the
10 HCl salt of α -Methyl-4-cyanophenylalanine ethyl ester (880 mg).

Step-3. To a solution of the compound obtained above in H₂O (30 mL) was added a solution of KH₂PO₄ (1.4 g) in H₂O (30 mL). To
15 this was added Lipase L (*Candida Lipolytica*, Sigma Aldrich, 1.4 g) and the pH of the suspension was adjusted to 6.40 using 1 N KOH. The progression of hydrolysis of ester to acid was monitored by HPLC (A = 0.1%TFA in H₂O, B = 0.1% TFA in MeCN; 15%B to 55%B over 20 minutes) with the acid eluting first (t =
20 5.8 minutes) followed by the ester (t = 10 minutes). The pH was maintained at 6.40 by the addition of further quantities of 1 N KOH until HPLC indicated the ratio of ester:acid equals 1:1.


After 31 hours, solid NaHCO₃ was added to bring the pH at 7.4
25 and the suspension was shaken with toluene (100 mL) and filtered through celite. The aqueous layer was separated and washed with DCM (2 x 200 mL) and the combined organics dried over MgSO₄. It was filtered and the filtrate was concentrated to furnish (R)- α -Methyl-4-cyanophenylalanine ethyl ester (340
30 mg).

Step 3: To a solution of the compound obtained above (340 mg) in DCM (10 mL) under N₂ at 0 °C was added neat 2,6-dichloro-4-

pyridyl isocyanate (305 mg). The reaction mixture was then warmed to room temperature and stirred for 4 hours whereupon it was concentrated *in vacuo* to give (R)-2-(4-cyanobenzyl)-5-(2,6-dichloro-4-pyridyl)-2-methylhydantoic acid ethyl ester (680 mg).

Step-4. After flushing with N₂, the compound obtained above was dissolved in dry EtOH (10 mL) and NaOEt (60 mg) was added. After stirring for 3 hours, water (10 mL) and EtOAc (10 mL) were added and the mixture was shaken. The aqueous phase was then separated and washed with EtOAc (3 x 10 mL) and the combined organics were dried over MgSO₄, filtered and concentrated *in vacuo*. Purification by liquid chromatography (EtOAc/Hexane 1/1) gave the titled compound (410 mg). MS (m/z) = 375 (M).


Example 5: 3-(2,6-Dichloro-4-pyridyl)-5-(4-Cyanobenzyl)-5-methyl-1-(5-carboxypentyl)-2,4-imidazolidinedione



The compound obtained in Example 4 (100 mg) was dissolved in a mixture of THF/MeOH (3 mL/1 mL). A solution of LiOH (25 mg in 1 mL H₂O) was added and the resulting mixture was stirred at room temperature for 5 hours. The pH of the mixture was adjusted to 3-4 using 1 M HCl and the mixture was extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The product was purified by

preparative TLC to give the titled compound (80 mg). MS: 489 (MH⁺).

**Example 6: (R)-3-(2,6-Dichloro-4-pyridyl)-5-(4-Cyanobenzyl)-
5 1,5-dimethyl-2,4-imidazolidinedione**



The compound obtained in Example 4 (340 mg) and KOtBu (132 mg) were weighed into a dry flask and flushed with N₂. The flask was placed in an ice bath and dry THF (9 mL) was added. After
10 stirring for 15 minutes, MeI (0.17 mL) was added and the reaction mixture was warmed to room temperature. After stirring for 1 hour, water (10 mL) and EtOAc (10 mL) were added and the mixture was shaken. The aqueous phase was then separated and washed with EtOAc (3 x 10 mL) and the combined
15 organics were dried over MgSO₄, filtered and concentrated in vacuo to give a white solid. Purification by liquid chromatography (EtOAc/Hexane = 1/1) gave the titled compound (260 mg). MS (m/z) = 389 (M).

20 The enantiomeric excess (e.e.) was determined to be >99% by chiral HPLC (0.5 mg/mL in MeOH, 3 μ L, Chiracel OD#ODOOCE-11030, 250 x 4.6 mm, isocratic gradient, Hexane/IPA).

**Example 7: (R)-3-(2,6-Dichloro-4-pyridyl)-5-(4-bromobenzyl)-5-
25 methyl-2,4-imidazolidinedione**

The titled compound was prepared in an analogous manner to that described for Example 4, using α -Methyl-4-bromophenylalanine ethyl ester which was prepared according to the methodology described in WO 98/39303. MS(m/z): 430 (MH).

5


Example 8: (R)-3-(2,6-Dichloro-4-pyridyl)-5-(4-bromobenzyl)-1,5-dimethyl-2,4-imidazolidinedione

The titled compound was prepared in an analogous manner to that described for Example 6. MS(m/z): 443 (MH).

The enantiomeric excess (e.e.) was determined to be >99% by chiral HPLC (0.5 mg/mL in MeOH, 3 μ L, Chiracel OD#ODOOCE-11030, 250 x 4.6 mm, isocratic gradient, Hexane/IPA).

15

Example 9: 3-(2,6-Dichloro-4-pyridyl)-5-(4-hydroxybenzyl)-5-methyl-2,4-imidazolidinedione



20 step-1. Et₃N (2.47 g) was added to a solution of the compound 1 (3 g) in H₂O (20 mL) and the resulting mixture was stirred

for 2 hours. The solution was extracted several times with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and evaporated to dryness. The resulting white solid (compound 2) was used as is without purification.


5

Step-2. A solution of 3,5-dichloro-4-pyridyl isocyanate (0.9 g) in CH₂Cl₂ (5 mL) was added to a solution of the compound obtained from step-1 (1 g) in anhydrous CH₂Cl₂ (15 mL) containing DMF (5 mL) at 0 °C. The reaction mixture was
10 allowed to warm to room temperature and stirred overnight. The mixture was partitioned between EtOAc/H₂O. The combined organic layers were dried over Na₂SO₄, filtered and concentrated to give 2-(4-hydroxybenzyl)-5-(2,6-dichloro-4-pyridyl)-2-methylhydantoic acid methyl ester (compound 3).
15 The product was used directly for the next step. MS: 398 (MH⁺).

Step-3. NaOEt (0.39 g) was added to a solution of the compound from step-2 (2.26 g) in anhydrous EtOH (15 mL) at 0 °C. The
20 yellow solution was then stirred at 0 °C for 5 hours and warmed to room temperature and stirred for 1 hour. EtOH was evaporated and the residue was partitioned between EtOAc/H₂O. The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The product was purified by flash
25 chromatography on silica gel (EtOAc/hexanes 1/1) to give the titled compound (1.4 g). MS: 366 (MH⁺).


Example 10: 3-(2,6-Dichloro-4-pyridyl)-5-(4-methoxybenzyl)-1,5-dimethyl-2,4-imidazolidinedione

30




The titled compound was obtained via methylation of the compound from Example 9 by following the procedure similar to Example 2. MS: 394 (MH^+).

Example 11: 3-(2,6-Dichloro-4-pyridyl)-5-(4-hydroxybenzyl)-1,5-dimethyl-2,4-imidazolidinedione



10 BBr_3 (1.14 mL, 1M in CH_2Cl_2) was added dropwise to a solution of the compound from Example 10 (0.15 g) in CH_2Cl_2 at 0 °C. The resulting mixture was stirred at 0 °C for 30 minutes and then stirred for an additional 30 minutes at room temperature. The reaction was quenched with water and partitioned between
 15 EtOAc and water. The aqueous solution was extracted with EtOAc and the combined organic layers were washed with water, dried over Na_2SO_4 , filtered and evaporated. The residue was purified via preparative TLC (EtOAc/Hexanes 1/1) to yield 0.125 g of the titled compound. MS: 380 (MH^+).

Example 12: 3-(2,6-Dichloro-4-pyridyl)-5-(4-i-propoxybenzyl)-1,5-dimethyl-2,4-imidazolidinedione




tBuOK (0.022g) was added to a solution of the compound from
 5 Example 11 (0.06 g) in THF (3 mL) and the solution was stirred
 for 5 minutes. 2-Iodopropane (0.054 g) was added and the
 reaction mixture was refluxed for 1.5 hours. The mixture was
 partitioned between EtOAc/water and the EtOAc layer was dried
 (Na₂SO₄), filtered and evaporated. The residue was purified
 10 via preparative TLC (EtOAc/Hexanes 1/1) to give the titled
 compound. MS: 422 (MH⁺).

The following compounds were prepared in a manner similar to
 Example 12.


15

Table 1



Example	R ²	Physicochemical Properties
13	CH ₃ (CH ₂) ₂ O-	MS: 422 (MH ⁺)
14	CH ₃ CH ₂ O-	MS: 408 (MH ⁺)

Example 15: 3-(2,6-Dichloro-4-pyridyl)-5-(4-ethoxy-3-fluorobenzyl)-1,5-dimethyl-2,4-imidazolidinedione.



5

To a solution of the compound from Example 14 (0.24 g) in CH₃CN (15 mL) was added 3,5-dichloro-1-fluoropyridinium triflate (0.38 g) and the mixture was refluxed for 30 hours. The mixture was concentrated and purified by HPLC to give the desired compound. MS m/z 426 (MH⁺).

Example 16: 3-(2,6-dichloro-4-pyridyl)-5-(4-(1,1,1trifluoromethoxybenzyl))-5-methyl-2,4-imidazolidinedione

15

The titled compound was prepared in a manner similar to Example 1. MS: 434 (MH⁺); mp. 151.2 °C.

Example 17: 3-(2,6-dichloro-4-pyridyl)-5-[4-(1,1,1trifluoromethoxybenzyl)]-1,5-Dimethyl-2,4-imidazolidinedione

The titled compound was obtained via methylation of the compound from Example 16 by following the procedure similar to Example 2. MS: 448 (MH⁺); mp. 113.7 °C.

Example 18: 3-(2,6-dichloro-4-pyridyl)-5-(4-fluorobenzyl)-5-methyl-2,4-imidazolidinedione

30 The titled compound was prepared in a manner similar to
Example 1. MS: 368 (MH⁺); mp 221.1 °C.

Example 19: 3-(2,6-dichloro-4-pyridyl)-5-(4-bromobenzyl)]-5-methyl-2,4-imidazolidinedione

5 The titled compound was prepared in a manner similar to Example 1. MS: 429 (MH⁺).

Example 20: 3-(2,6-dichloro-4-pyridyl)-5-(4-bromobenzyl)]-1,5-dimethyl-2,4-imidazolidinedione

10

The titled compound was obtained via methylation of the compound from Example 19 by following the procedure similar to Example 2. MS: 443 (MH⁺).


15 **Cell Adhesion Protocol**

Cell Adhesion The recombinant protein ICAM-1•Fc was constructed from the 5 extracellular domains of human ICAM-1 and fusion with the constant region of human IgG. ICAM-1•Fc was purified by Protein A affinity chromatography and stored
20 in aliquots at -20°C. Immobilized ICAM-1•Fc was prepared by dilution of the protein in PBS pH 7.5, transfer of 100 µl/well to Falcon Probind III plates and overnight incubation at 4°C. Wells coated with BSA served as a measure of non-specific background adhesion. Washed plates were blocked with a
25 solution of 0.25% ovalbumin in PBS for 1 h at 37°C. HBSS washed Jurkat cells were suspended to a final concentration of 2.5x10⁶/ml in TBSg adhesion buffer (24 mM Tris pH 7.4, 0.14 M NaCl, 2.7 mM KCl, 2 mM glucose, 0.1% HSA. A 100 µl volume of cells was added to the blocked and washed ICAM-1•Fc coated
30 plates that contained 100 µl of plate buffer (TBSg, 10 mM MgCl₂, 2% DMSO). Adhesion was for 1 h at 37°C. Non-adherent cells were removed using the EL404 plate washer (BioTek Instruments; Highland Park, VT). The number of adherent cells was quantified by measuring enzymatic activity of endogenous

N-acetyl-hexosaminidase using the enzyme substrate *p*-nitrophenol-N-acetyl- β -D-glucosaminide, pNAG. The amount of liberated *p*-nitrophenol was measured by reading the optical density at 405 nm using a vertical pathway spectrophotometer
5 to quantify cell attachment (VMAX Kinetic Microplate Reader, Molecular Devices, Menlo Park, CA). For competition studies the compounds from 100% DMSO stock solutions were diluted in plate buffer at 2-fold the required testing concentration prior to transfer to the ICAM-1-Fc coated plate and serial
10 dilution.

Claims

1. A compound of the formula (I):



5 or a pharmaceutically acceptable salt thereof, wherein
 R¹ is selected from

- 1) hydrogen atom, or
 - 2) a C₁₋₆ alkyl group which may be optionally substituted
 with carboxyl group or a C₁₋₆ alkoxy carbonyl group;
- 10 R^{2a} and R^{2b} are independently hydrogen atom, a halogen atom,
 hydroxyl group, cyano group, a C₁₋₆ alkyl group which may be
 optionally substituted with 1 to 3 halogen atoms, a C₁₋₆
 alkylthio group which may be optionally substituted with 1 to
 3 halogen atoms, a C₁₋₆ alkylsulfinyl group which may be
 15 optionally substituted with 1 to 3 halogen atoms, a C₁₋₆
 alkylsulfonyl group which may be optionally substituted with 1
 to 3 halogen atoms, or a C₁₋₆ alkoxy group which may be
 optionally substituted with 1 to 3 halogen atoms;
 R³ is a C₁₋₆ alkyl group; and
- 20 R⁴ and R⁵ are independently a halogen atom.

2. The compound according to claim 1, wherein R¹ is hydrogen
 atom or a C₁₋₆ alkyl group which may be optionally substituted
 with carboxyl group or a C₁₋₆ alkoxy carbonyl group, R^{2a} and R^{2b}
 25 are independently hydrogen atom, a halogen atom, hydroxyl
 group, cyano group or a C₁₋₆ alkoxy group which may be

optionally substituted with 1-3 halogen atoms, R³ is a C₁₋₆ alkyl group, and R⁴ and R⁵ are independently a halogen atom.

3. The compound according to claim 2, wherein R¹ is hydrogen atom or a C₁₋₆ alkyl group, one of R^{2a} and R^{2b} is hydrogen atom, and the other is a halogen atom, cyano group, or a C₁₋₆ alkoxy group which may be optionally substituted with 1-3 halogen atoms, R³ is a C₁₋₆ alkyl group, and R⁴ and R⁵ are independently a halogen atom.

10

4. The compound according to claim 3, wherein R¹ is hydrogen atom or methyl group, one of R^{2a} and R^{2b} is hydrogen atom and the other is bromine atom, cyano group, a C₁₋₆ alkoxy group or trifluoromethoxy group, R³ is methyl group, R⁴ and R⁵ are chlorine atom.


5. The compound according to claim 2, wherein R¹ is a C₁₋₆ alkyl group which is substituted with a C₁₋₆ alkoxycarbonyl group or carboxyl group, one of R^{2a} and R^{2b} is hydrogen atom, and the other is a halogen atom, cyano group, or C₁₋₆ alkoxy group which may be optionally substituted with 1-3 halogen atoms, R³ is a C₁₋₆ alkyl group, and R⁴ and R⁵ are independently a halogen atom.

6. The compound according to claim 5, R³ is methyl group, and R⁴ and R⁵ are chlorine atom.

7. The compound according to claim 1, wherein the compound is selected from:
3-(2,6-Dichloro-4-pyridyl)-5-(4-bromobenzyl)-1,5-dimethyl-2,4-imidazolidinedione,
3-(2,6-Dichloro-4-pyridyl)-5-(4-propoxybenzyl)-1,5-dimethyl-2,4-imidazolidinedione,

3- (2,6-Dichloro-4-pyridyl) -5- (4-ethoxybenzyl) -1,5-dimethyl-
 2,4-imidazolidinedione,
 3- (2,6-dichloro-4-pyridyl) -5- (4-
 (1,1,1trifluoromethoxybenzyl)] -5-methyl-2,4-
 5 imidazolidinedione,
 3- (2,6-dichloro-4-pyridyl) -5- [4-
 (1,1,1trifluoromethoxybenzyl)] -1,5-Dimethyl-2,4-
 imidazolidinedione,
 3- (2,6-Dichloro-4-pyridyl) -5- (4-Cyanobenzyl) -5-methyl-2,4-
 10 imidazolidinedione,
 3- (2,6-Dichloro-4-pyridyl) -5- (4-Cyanobenzyl) -1,5-dimethyl-2,4-
 imidazolidinedione (TR-15170); and
 a pharmaceutically acceptable salt of these compounds.


15 7. A process for preparing a compound of the formula (I-a):



or a pharmaceutically acceptable salt thereof, wherein
 R^{2a} and R^{2b} are independently hydrogen atom, a halogen atom,
 hydroxyl group, cyano group, a C_{1-6} alkyl group which may be
 20 optionally substituted with 1 to 3 halogen atoms, a C_{1-6}
 alkylthio group which may be optionally substituted with 1 to
 3 halogen atoms, a C_{1-6} alkylsulfinyl group which may be
 optionally substituted with 1 to 3 halogen atoms, a C_{1-6}
 alkylsulfonyl group which may be optionally substituted with 1
 25 to 3 halogen atoms, or a C_{1-6} alkoxy group which may be
 optionally substituted with 1 to 3 halogen atoms;
 R^3 is a C_{1-6} alkyl group;


R^4 and R^5 are independently a halogen atom;

which comprises cyclizing the compound of the formula (II):



wherein OR^6 is a hydroxyl group or a protected hydroxyl group,
 5 and the other symbols are the same as defined above, and
 converting into the pharmaceutically acceptable salt, if
 necessary.

8. A process for preparing a compound of the formula (I-b):



10


or a pharmaceutically acceptable salt thereof, wherein
 R^{11} is a C_{1-6} alkyl group which may be optionally substituted
 with carboxyl group or a C_{1-6} alkoxy carbonyl group;
 R^{2a} and R^{2b} are independently hydrogen atom, a halogen atom,
 15 hydroxyl group, cyano group, a C_{1-6} alkyl group which may be
 optionally substituted with 1 to 3 halogen atoms, a C_{1-6}
 alkylthio group which may be optionally substituted with 1 to
 3 halogen atoms, a C_{1-6} alkylsulfinyl group which may be
 optionally substituted with 1 to 3 halogen atoms, a C_{1-6}
 20 alkylsulfonyl group which may be optionally substituted with 1

to 3 halogen atoms, or a C₁₋₆ alkoxy group which may be optionally substituted with 1 to 3 halogen atoms;

R³ is a C₁₋₆ alkyl group;

R⁴ and R⁵ are independently a halogen atom;


5 which comprises (1) alkylating a compound of the formula (I-a):



wherein the symbols are the same as defined above,

10 (2) hydrolyzing the resulting compound, if necessary, and
(3) converting into the pharmaceutically acceptable salt, if further desired.

8. A process for preparing a compound of the formula (I-b):



or a pharmaceutically acceptable salt thereof, wherein


R¹¹ is a C₁₋₆ alkyl group which may be optionally substituted with carboxyl group or a C₁₋₆ alkoxycarbonyl group;

R²¹ is a C₁₋₆ alkoxy group which may be optionally substituted
20 with 1 to 3 halogen atoms;

R^3 is a C_{1-6} alkyl group;

R^4 and R^5 are independently a halogen atom;

which comprises alkylating a compound of the formula (I-d):



5 wherein R^1 is hydrogen atom or a C_{1-6} alkyl group which may be optionally substituted with carboxyl group or a C_{1-6} alkoxy carbonyl group, and the other symbols are the same as defined above, and converting into the pharmaceutically acceptable salt, if necessary.

10

9. A pharmaceutical composition which comprises a therapeutically effective amount of the compound as set forth in any one of claims 1-6 in admixture with a therapeutically acceptable carrier or diluent.

15

10. A method for treatment or prevention of $\alpha_L\beta_2$ adhesion mediated condition in a mammal comprising administering a therapeutically effective amount of the compound as set forth in any one of claims 1-6.

20

11. The method according to claim 10, said condition is selected from rheumatoid arthritis, asthma, allergy conditions, adult respiratory distress syndrome, AIDS, cardiovascular diseases, thrombosis, harmful platelet aggregation, reocclusion following thrombolysis, reperfusion injury, skin inflammatory diseases, osteoporosis, osteoarthritis, arteriosclerosis, neoplastic diseases, wound,

25

detaching retina, Type I diabetes, multiple sclerosis, systemic lupus erythematosus, ophthalmic inflammatory conditions, inflammatory bowel diseases, regional enteritis, Sjogren's Syndrome, and rejection after transplantation.

5

12. The method according to claim 11, wherein said diseases is selected from psoriasis, rheumatoid arthritis, inflammatory bowel diseases (Crohn's disease, ulcerative colitis), systemic lupus erythematosus, atopic dermatitis, Sjogren's Syndrome, rejection after transplantation (allograft rejection and graft vs. host disease).

10